arXiv:1512.00992v2 [math.AG] 7 Dec 2015

# ULRICH BUNDLES ON ABELIAN SURFACES

ARNAUD BEAUVILLE

ABSTRACT. We prove that any abelian surface admits a rank 2 Ulrich bundle.

Let $X \subset \mathbb{P}^N$ be a projective variety of dimension $d$ over an algebraically closed field. An *Ulrich bundle* on $X$ is a vector bundle $E$ on $X$ satisfying $H^*(X, E(-1)) = \ldots = H^*(X, E(-d)) = 0$. This notion was introduced in [ES], where various other characterizations are given; let us just mention that it is equivalent to say that $E$ admits a linear resolution as a $\mathscr{O}_{\mathbb{P}^N}$-module, or that the pushforward of $E$ onto $\mathbb{P}^d$ by a general linear projection is a trivial bundle.

In [ES] the authors ask whether every projective variety admits an Ulrich bundle. The answer is known only in a few cases: hypersurfaces and complete intersections [HUB], del Pezzo surfaces [ES, Corollary 6.5]. The case of K3 surfaces is treated in [AFO]. In this short note we show that the existence of Ulrich bundles for abelian surfaces follows easily from Serre's construction:

**Theorem.** *Any abelian surface $X \subset \mathbb{P}^N$ carries a rank 2 Ulrich bundle.*

*Proof* : We put $\dim H^0(X, \mathscr{O}_X(1)) = n$. Let $C$ be a smooth curve in $|\mathscr{O}_X(1)|$; we have $\mathscr{O}_C(1) \cong \omega_C$, and $g(C) = n+1$. We choose a subset $Z \subset C$ of $n$ general points. Then $Z$ has the *Cayley-Bacharach property* on $X$ (see for instance [HL], Theorem 5.1.1): for every $p \in Z$, any section of $H^0(X, \mathscr{O}_X(1))$ vanishing on $Z \smallsetminus \{p\}$ vanishes on $Z$. Indeed, the image $V$ of the restriction map $H^0(X, \mathscr{O}_X(1)) \to H^0(C, \mathscr{O}_C(1))$ has dimension $n-1$, hence the only element of $V$ vanishing on $n-1$ general points is zero; thus the only element of $|\mathscr{O}_X(1)|$ containing $Z \smallsetminus \{p\}$ is $C$.

By *loc. cit.*, there exists a rank 2 vector bundle $E$ on $X$ and an exact sequence

$$0 \to \mathscr{O}_X \xrightarrow{s} E \longrightarrow \mathscr{I}_Z(1) \to 0\,.$$

Let $\eta$ be a nonzero element of $\mathrm{Pic}^o(X)$; then $h^0(\omega_C \otimes \eta) = n$, so $H^0(C, \omega_C \otimes \eta(-Z)) = 0$ since $Z$ is general, and therefore $H^0(X, \mathscr{I}_Z\eta(1)) = 0$. Since $\chi(\mathscr{I}_Z\eta(1))) = 0$ we have also $H^1(X, \mathscr{I}_Z\eta(1)) = 0$; from the above exact sequence we conclude that $H^*(X, E \otimes \eta) = 0$.

The zero locus of the section $s$ of $E$ is $Z$; since $\det E = \omega_C$, we get an exact sequence

$$0 \to \mathscr{O}_C(Z) \xrightarrow{s_{|C}} E_{|C} \longrightarrow \omega_C(-Z) \to 0\,.$$

As above the cohomology of $\omega_C \otimes \eta(-Z)$ and $\eta(Z)$ vanishes, hence $H^*(C, (E \otimes \eta)_{|C}) = 0$. Now from the exact sequence

$$0 \to E(-1) \to E \to E_{|C} \to 0$$

we conclude that $H^*(X, E \otimes \eta(-1)) = H^*(X, E \otimes \eta) = 0$, hence $E \otimes \eta(1)$ is an Ulrich bundle. ∎

I am indebted to M. Aprodu for pointing out an inaccuracy in the first version of this note.

## REFERENCES

[AFO] M. Aprodu, G. Farkas, A. Ortega : *Minimal resolutions, Chow forms and Ulrich bundles on $K3$ surfaces*. Preprint `arXiv:1212.6248`. J. Reine Angew. Math., to appear.

[HUB] J. Herzog, B. Ulrich, J. Backelin : *Linear maximal Cohen-Macaulay modules over strict complete intersections*. J. Pure Appl. Algebra **71** (1991), no. 2-3, 187-202.

[ES] D. Eisenbud, F.-O. Schreyer : *Resultants and Chow forms via exterior syzygies*. J. Amer. Math. Soc. **16** (2003), no. 3, 537-579.

[HL] D. Huybrechts, M. Lehn : *The geometry of moduli spaces of sheaves* (2nd edition). Cambridge University Press, Cambridge, 2010.

LABORATOIRE J.-A. DIEUDONNÉ, UMR 7351 DU CNRS, UNIVERSITÉ DE NICE, PARC VALROSE, F-06108 NICE CEDEX 2, FRANCE

*E-mail address*: `arnaud.beauville@unice.fr`